

# ASSIGNMENT 2

## L<sup>A</sup>T<sub>E</sub>XPresentation

Ritika Kamboj  
MAT/20/122  
20044563047

Mata Sundri College For Women  
University of Delhi

# Content of the page number 69

- Let  $\mathbf{x} = x_1, \dots, x_n$ , where the  $x_i$  are non-negative real numbers. Set

$$M_r(\mathbf{x}) = \left( \frac{x_1^r + x_2^r + \cdots + x_n^r}{n} \right)^{1/r}, \quad r \in \mathbf{R} \setminus \{0\},$$

and

$$M_0(\mathbf{x}) = (x_1 x_2 \cdots x_n)^{1/n}.$$

We call  $M_r(\mathbf{x})$  the  $r^{th}$  power mean of  $\mathbf{x}$ .

*Claim:*

$$\lim_{r \rightarrow 0} M_r(\mathbf{x}) = M_0(\mathbf{x})$$

- Define

$$V_n = \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ x_1 & x_2 & x_3 & \cdots & x_n \\ x_1^2 & x_2^2 & x_3^2 & \cdots & x_n^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_1^{n-1} & x_2^{n-1} & x_3^{n-1} & \cdots & x_n^{n-1} \end{bmatrix}$$

We call  $V_n$  the *Vandermonde matrix* of order n.

Claim:

$$\det V_n = \prod_{1 \leq i < j \leq n} (x_j - x_i)$$

## Question. 4



$$3^3 + 4^3 + 5^3 = 6^3$$



$$\sqrt{100} = 10$$



$$(a+b)^3 = a^3 + b^3 + 3a^2b + 3ab^2$$



$$\sum_{k=1}^n k = \frac{n(n+1)}{2}$$



$$\frac{\pi}{4} = \frac{1}{1} - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \frac{1}{11} + \dots$$

## Remaining parts of Ques.4



$$\cos \theta = \sin(90^\circ - \theta)$$



$$e^{i\theta} = \cos \theta + i \sin \theta$$



$$\lim_{x \rightarrow \infty} \frac{\sin \theta}{\theta} = 1$$



$$\lim_{x \rightarrow \infty} \frac{\pi(x)}{x / \log x} = 1$$



$$\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}$$

## Question 5

- Positive numbers  $a, b$  and  $c$  are the side lengths of a triangle if and only if  $a + b > c, b + c > a$ , and  $c + a > b$ .
- The area of a triangle with side lengths  $a, b, c$  is given by *Heron's formula*:

$$A = \sqrt{s(s - a)(s - b)(s - c)},$$

where  $s$  is the semiperimeter  $(a+b+c)/2$ .

- The volume of a regular tetrahedron of edge length 1 is  $\sqrt{2}/12$ .
- The quadratic equation  $ax^2 + bx + c = 0$  has roots

$$r_1, r_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## Remaining Parts of Ques.5

- The *derivative* of a function  $f$ , denoted  $f'$ , is defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}.$$

- A real-valued function  $f$  is *convex* on an interval  $I$  if

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y),$$

for all  $x, y \in I$  and  $0 \leq \lambda \leq 1$

- The general solution to the differential equation

$$y'' - 3y' + 2y = 0$$

is

$$y = C_1 e^x + C_2 e^{2x}.$$

## Remaining part of Ques.5

- The *Fermat number*  $F_n$  is defined as

$$F_n = 2^{2^n}, n \geq 0.$$

## Question 6



$$\frac{d}{dx} \left( \frac{x}{x+1} \right) = \frac{1}{(x+1)^2}$$



$$\lim_{n \rightarrow \infty} \left( 1 + \frac{1}{n} \right)^n = e$$



$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$



$$R_\theta = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

## Remaining Parts of Ques.6



$$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \mathbf{k}$$



$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{bmatrix}$$



$$f(x) = \begin{cases} -x^2, & x < 0 \\ x^2, & 0 \leq x \leq 2 \\ 4, & x > 2 \end{cases} .$$

## Question 7(1)

$$1 + 2 = 3$$

$$4 + 5 + 6 = 7 + 8$$

$$9 + 10 + 11 + 12 = 13 + 14 + 15$$

$$16 + 17 + 18 + 19 + 20 = 21 + 22 + 23 + 24$$

$$25 + 26 + 27 + 28 + 29 + 30 = 31 + 32 + 33 + 34 + 35$$

## Question 7(2)

$$\begin{aligned}(a + b)^2 &= (a + b)(a + b) \\&= (a + b)a + (a + b)b \\&= a(a + b) + b(a + b) \\&= a^2 + ab + ba + b^2 \\&= a^2 + ab + ab + b^2 \\&= a^2 + 2ab + b^2\end{aligned}$$

## Question 7(3)

$$\begin{aligned}\tan(\alpha + \beta + \gamma) &= \frac{\tan(\alpha + \beta) + \tan \gamma}{1 - \tan(\alpha + \beta) \tan \gamma} \\&= \frac{\frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} + \tan \gamma}{1 - \left( \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} \right) \tan \gamma} \\&= \frac{\tan \alpha + \tan \beta + (1 - \tan \alpha \tan \beta) \tan \gamma}{1 - \tan \alpha \tan \beta - (\tan \alpha + \tan \beta) \tan \gamma} \\&= \frac{\tan \alpha + \tan \beta + \tan \gamma - \tan \alpha \tan \beta \tan \gamma}{1 - \tan \alpha \tan \beta - \tan \alpha \tan \gamma - \tan \beta \tan \gamma}\end{aligned}$$

## Question 7(4)

$$\begin{aligned}\prod_p \left(1 - \frac{1}{p^2}\right) &= \prod_p \frac{1}{1 + \frac{1}{p^2} + \frac{1}{p^4} + \dots} \\&= \left( \prod_p \left(1 + \frac{1}{p^2} + \frac{1}{p^4} + \dots\right) \right)^{-1} \\&= \left(1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots\right)^{-1} \\&= \frac{6}{\pi^2}\end{aligned}$$

