


ASSIGNMENT-1

Q1). Graph each of the functions. Experiment with different domains or viewpoints to display the best images.


(A) $f(x) = x/(1+x^2)$

In[1]:= `Plot[x/(1+x^2), {x, -5, 5}]`




(B) $y = x \sin(1/x)$

In[3]:= `Plot[x Sin[1/x], {x, -Pi, Pi}]`




(C) $g(x,y) = \cos(x) + \sin(y)$

In[12]:= `Plot3D[Cos[x]+Sin[y], {x, -Pi, Pi}, {y, -Pi, Pi}]`



(D) $z = \frac{xy}{x^2 + y^2}$

In[4]:= `Plot3D[{x y / (x^2 + y^2)}, {x, -5, 5}, {y, -5, 5}]`



Q2). Let $f(x) = \frac{x}{1+x^2}$

In[20]:= `f[x_] := x / (1 + x^2)`

(A) Find $f'(x)$ and $f''(x)$

In[28]:= **D[f[x], x]**

$$\text{Out[28]} = -\frac{2x^2}{(1+x^2)^2} + \frac{1}{1+x^2}$$

In[23]:= **D[f[x], {x, 2}]**

$$\text{Out[23]} = -\frac{4x}{(1+x^2)^2} + x \left(\frac{8x^2}{(1+x^2)^3} - \frac{2}{(1+x^2)^2} \right)$$

(B) Find $f'(-1)$ and $f'(0)$

In[24]:= **f'[-1]**

Out[24]= 0

In[25]:= **f'[0]**

Out[25]= 1

(C) Find $f''(0)$ and $f''(1)$

In[26]:= **f''[0]**

Out[26]= 0

In[27]:= **f''[1]**

$$\text{Out[27]} = -\frac{1}{2}$$

Q3). Find the prime factorization of each integer:

(A) 3527218133309949276293

In[30]:= **FactorInteger [3 527 218 133 309 949 276 293]**

Out[30]= {{15 013 , 2}, {25 013 , 3}}

(B) 471945325930166269

In[31]:= **FactorInteger [471 945 325 930 166 269]**

Out[31]= {{4211 , 1}, {34 589 , 1}, {46 747 , 1}, {69 313 , 1}}

(C) 471945325930166281

In[32]:= **FactorInteger** [471 945 325 930 166 281]

Out[32]= {{471 945 325 930 166 281 , 1}}

Q4). Complete each expression. Do you notice a pattern?

(A) $3^6 \bmod 7$

In[33]:= **Mod**[3 ^ 6, 7]

Out[33]= 1

(B) $6^{10} \bmod 11$

In[34]:= **Mod**[6 ^ 10, 11]

Out[34]= 1

(C) $7^{20} \bmod 21$

In[35]:= **Mod**[7 ^ 20, 21]

Out[35]= 7

(D) $7^{22} \bmod 23$

In[36]:= **Mod**[7 ^ 22, 23]

Out[36]= 1

Q8). Let $M = \begin{Bmatrix} 1 & 1 \\ 1 & 0 \end{Bmatrix}$

In[38]:= **M** = {{1, 1}, {1, 0}}

Out[38]= {{1, 1}, {1, 0}}

(A) Find M^2, M^3, \dots, M^{10}

In[39]:= **MatrixPower** [M, 2]

Out[39]= {{2, 1}, {1, 1}}

In[40]:= **MatrixPower** [M, 3]

Out[40]= {{3, 2}, {2, 1}}

In[41]:= **MatrixPower** [M, 4]

Out[41]= {{5, 3}, {3, 2}}

In[42]:= **MatrixPower [M, 5]**

Out[42]= {{8, 5}, {5, 3}}

In[43]:= **MatrixPower [M, 6]**

Out[43]= {{13, 8}, {8, 5}}

In[44]:= **MatrixPower [M, 7]**

Out[44]= {{21, 13}, {13, 8}}

In[45]:= **MatrixPower [M, 8]**

Out[45]= {{34, 21}, {21, 13}}

In[46]:= **MatrixPower [M, 9]**

Out[46]= {{55, 34}, {34, 21}}

In[47]:= **MatrixPower [M, 10]**

Out[47]= {{89, 55}, {55, 34}}

(B) Do your answers suggest a way to compute Fibonacci numbers? Find the 100th Fibonacci number.

In[48]:= **f[0] = 1;**

In[49]:= **f[1] = 1;**

In[50]:= **f[n_] := f[n] = f[n - 2] + f[n - 1]**

In[51]:= **f[100]**

Out[51]= 573 147 844 013 817 084 101

Q9). Find solutions to the following equations or systems of equations:

(A) Find x, if $x^2+x=1$

In[54]:= **Solve[{x^2 + x == 1}, x]**

Out[54]= $\left\{ \left\{ x \rightarrow \frac{1}{2}(-1 - \sqrt{5}) \right\}, \left\{ x \rightarrow \frac{1}{2}(-1 + \sqrt{5}) \right\} \right\}$

(B) Find x, if $x^2+x=-1$

In[55]:= **Solve[{x^2 + x == -1}, x]**

Out[55]= $\left\{ \left\{ x \rightarrow -(-1)^{1/3} \right\}, \left\{ x \rightarrow (-1)^{2/3} \right\} \right\}$

(C) Find x and y

$$4x-3y=5$$

$$6x+2y=14$$

In[56]:= `Solve[{4 x - 3 y == 5 && 6 x + 2 y == 14}, {x, y}]`

Out[56]= `{{x -> 2, y -> 1}}`

(D) Find x,y,z and t

$$-2x-2y+3z+t=8$$

$$-3x+0y-6z+t=-19$$

$$6x-8y+6z+5t=47$$

$$x+3y-3z-t=-9$$

In[57]:= `Solve[{-2 x - 2 y + 3 z + t == 8 && -3 x + 0 y - 6 z + t == -19 && 6 x - 8 y + 6 z + 5 t == 47 && x + 3 y - 3 z - t == -9}, {x, y, z, t}]`

Out[57]= `{{x -> 2, y -> 1, z -> 3, t -> 5}}`

Q10). Assume that I invest \$250 at the beginning of the year, \$300 at the beginning of the second quarter, \$350 at the beginning of the third quarter, and \$400 at the beginning of the fourth quarter. At the end of the year, I have \$1365 (because my investments grow). To find my (continuous) rate of return, solve this equation for r :

$$250e^{1.0r} + 300e^{0.75r} + 350e^{0.5r} + 400e^{0.25r} = 1365$$

In[65]:= `FindRoot[{250 e^(1.0 r) + 300 e^(0.75 r) + 350 e^(0.5 r) + 400 e^(0.25 r) == 1365}, {r, 0}]`

Out[65]= `{r -> 0.084104}`

Q11). If n is a positive number, and $g > 0$ is any "guess" for the square root of n, then a better estimate of \sqrt{n} is the average of g and n/g , i.e., $(g+(n/g))/2$. Write a function called mysqrt that accepts one argument, begins with an initial guess of 1.0, finds 20 new guesses, returns the answer.

In[66]:= `mysqrt[n_] := Module[{i = 1, g = 1}, While[i ≤ 20, g = (g + (n/g))/2; i = i + 1]; g]`

In[97]:= `N[mysqrt[2], 6]`

Out[97]= 1.41421

In[98]:= `N[Sqrt[2], 6]`

Out[98]= 1.41421

```
In[99]:= N[mysqrt[3]]
Out[99]= 1.73205
```

Q12). The Collatz conjecture states that if we start from any natural number $a_0=n$ and form a sequence by the rule

$$a_{i+1} = \begin{cases} a_i/2, & \text{if } a_i \text{ is even} \\ 3a_i+1, & \text{if } a_i \text{ is odd} \end{cases}$$

then the sequence eventually contains the value 1. For example, starting from $a_0=6$, we get the sequence 6, 3, 10, 5, 16, 8, 4, 2, 1 (we reached 1 after eight steps).

(A) Write a (recursive) function called `collatz` that accepts a single argument, n , and returns :

- 0 if n is equal to 1
- $1+\text{collatz}(n/2)$ if n is even
- $1+\text{collatz}(3*n+1)$ if n is odd

Thus, $\text{collatz}(n)$ is the number of steps needed to go from n to 1

```
In[100]:= Clear[collatz];
In[101]:= collatz[n_] := Which[n == 1, collatz[n] = 0, EvenQ[n],
    collatz[n] = 1 + collatz[n/2], OddQ[n], collatz[n] = 1 + collatz[3 * n + 1]];

```

(B) Verify the values :

```
n      : collatz(n)
1      :      0
2      :      1
6      :      8
27     :     111
```

```
In[103]:= collatz[1]
Out[103]= 0
```

```
In[104]:= collatz[2]
Out[104]= 1
```

```
In[105]:= collatz[6]
Out[105]= 8
```

```
In[106]:= collatz[27]
Out[106]= 111
```