

PRACTICAL NO.1

PRAKRITI RATURI


MAT/19/95

Que1. Graph each of the functions. Experiment with different domains or view points to display the best images.

a) $f(x) = x/(1+x^2)$


```
In[1]:= f[x_] := x / (1 + x^2)
```

```
In[4]:= Plot[f[x], {x, -10, 10}]
```



b) $y = x \sin(1/x)$


```
In[5]:= Plot[x * Sin[1/x], {x, -2, 2}]
```



c) $g(x,y) = \cos(x) + \sin(y)$

```
In[13]:= g[x_, y_] := Cos[x] + Sin[y]
```


```
In[15]:= Plot3D[g[x, y], {x, 0, 2 Pi}, {y, 0, 2 Pi}]
```



d) $z = xy/(x^2+y^2)$

```
In[16]:= z[x_, y_] := x * y / (x^2 + y^2)
```

In[18]:= Plot3D[z[x, y], {x, -5, 5}, {y, -5, 5}]



Que2. $f(x) = x/(1+x^2)$

In[32]:= $f[x_] := x / (1 + x^2)$

In[33]:= $A = D[f[x], x]$

$$\text{Out}[33]= -\frac{2 x^2}{(1+x^2)^2} + \frac{1}{1+x^2}$$

In[34]:= $B = D[f'[x], x]$

$$\text{Out}[34]= \frac{8 x^3}{(1+x^2)^3} - \frac{6 x}{(1+x^2)^2}$$

In[36]:= $A /. \{x \rightarrow -1\}$

$$\text{Out}[36]= 0$$

In[38]:= $A /. \{x \rightarrow 0\}$

$$\text{Out}[38]= 1$$

In[39]:= $B /. \{x \rightarrow 0\}$

$$\text{Out}[39]= 0$$

In[40]:= $B /. \{x \rightarrow 1\}$

$$\text{Out}[40]= -\frac{1}{2}$$

Que3.

a) 3,527,218,133,309,949,276,293

```
In[41]:= FactorInteger [3 527 218 133 309 949 276 293 ]
Out[41]= {{15 013 , 2}, {25 013 , 3}}
```

b) 471,945,325,930,166,269

```
In[42]:= FactorInteger [471 945 325 930 166 269 ]
Out[42]= {{4211 , 1}, {34 589 , 1}, {46 747 , 1}, {69 313 , 1}}
```

c) 471,945,325,930,166,281

```
In[43]:= FactorInteger [471 945 325 930 166 281 ]
Out[43]= {{471 945 325 930 166 281 , 1}}
```

Que4.

a) $3^6 \bmod 7$

```
In[44]:= Mod[3 ^ 6, 7]
Out[44]= 1
```

b) $6^{10} \bmod 11$

```
In[45]:= Mod[6 ^ 10, 11]
Out[45]= 1
```

c) $7^{20} \bmod 21$

```
In[46]:= Mod[7 ^ 20, 21]
Out[46]= 7
```

d) $7^{22} \bmod 23$

```
In[47]:= Mod[7 ^ 22, 23]
Out[47]= 1
```

Que8.

a)

```
In[1]:= M = {{1, 1}, {1, 0}}
Out[1]= {{1, 1}, {1, 0}}

In[2]:= MatrixPower [M, 2]
Out[2]= {{2, 1}, {1, 1}}

In[3]:= MatrixPower [M, 3]
Out[3]= {{3, 2}, {2, 1}}

In[4]:= MatrixPower [M, 4]
Out[4]= {{5, 3}, {3, 2}}

In[5]:= MatrixPower [M, 5]
Out[5]= {{8, 5}, {5, 3}}

In[6]:= MatrixPower [M, 6]
Out[6]= {{13, 8}, {8, 5}}

In[7]:= MatrixPower [M, 7]
Out[7]= {{21, 13}, {13, 8}}

In[8]:= MatrixPower [M, 8]
Out[8]= {{34, 21}, {21, 13}}

In[9]:= MatrixPower [M, 9]
Out[9]= {{55, 34}, {34, 21}}

In[10]:= MatrixPower [M, 10]
Out[10]= {{89, 55}, {55, 34}}
```

b)

```
In[78]:= ClearAll[f]
```

```
In[79]:= f[0] = 1;
In[80]:= f[1] = 1;
In[81]:= f[n_] := f[n] = f[n - 2] + f[n - 1];
In[82]:= f[n] /. {n → 100}
Out[82]= 573 147 844 013 817 084 101
```

Que9.

a)

```
In[90]:= Solve[x^2 + x == 1, x]
Out[90]= {{x → 1/2 (-1 - √5)}, {x → 1/2 (-1 + √5)}}
```

b)

```
In[91]:= Solve[x^2 + x == -1, x]
Out[91]= {{x → -(-1)^{1/3}}, {x → (-1)^{2/3}}}
```

c)

```
In[92]:= Solve[{4 x - 3 y == 5, 6 x + 2 y == 14}, {x, y}]
Out[92]= {{x → 2, y → 1}}
```

d)

```
In[93]:= Solve[{-2 x - 2 y + 3 z + t == 8, -3 x + 0 y - 6 z + t == -19,
6 x - 8 y + 6 z + 5 t == 47, x + 3 y - 3 z - t == -9}, {x, y, z, t}]
Out[93]= {{x → 2, y → 1, z → 3, t → 5}}
```

Que10.

```
In[1]:= FindRoot[
250 * Exp[1.0 * r] + 300 * Exp[0.75 * r] + 350 * Exp[0.5 * r] + 400 * Exp[0.25 * r] == 1365, {r, 0}]
Out[1]= {r → 0.084104}
```

Que11.

```
In[2]:= mysqrt[n_] := Module[{i = 1, g = 1}, While[i <= 20, g = (g + n/g)/2; i = i + 1]; g]
In[3]:= N[mysqrt[2], 6]
Out[3]= 1.41421
In[4]:= N[mysqrt[3]]
Out[4]= 1.73205
```

Que12.

```
In[7]:= Clear[collatz];
In[8]:= collatz[n_] := Which[n == 1, collatz[n] = 0, EvenQ[n],
                           collatz[n] = 1 + collatz[n/2], OddQ[n], collatz[n] = 1 + collatz[3*n + 1]]
In[9]:= collatz[27]
Out[9]= 111
```